

RESTRICTED USE PESTICIDE

Due to acute oral and inhalation toxicity.
For retail sale and application only by certified applicators or personnel under their direct supervision.

EPA SLN NO.
ID-920002



FOR DISTRIBUTION
AND USE ONLY IN
IDAHO

Furadan® 4F Insecticide-Nematicide

For Agricultural or Commercial Use Only

EPA REG. No. 279-2876 ZA

EMERGENCY CALLS: 800-331-3148

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER
INCONSISTENT WITH ITS LABELING.

CROP	PEST	RATE OF APPLICATION
Sugarbeets	Sugarbeet Root Maggot	1 to 2 quarts per acre

DIRECTIONS FOR USE:

Make application in a 4 to 5 inch band over the plant row. Apply in a minimum of 10 gallons of finished spray per acre. Time treatments so that applications begin after adult fly emergence. Under sprinkler irrigation follow treatment with a light irrigation. Moisture after treatment is necessary to incorporate and activate the Furadan 4F. Rainfall will be needed in furrow irrigation fields to achieve consistent results. Fungicides, herbicides, or liquid fertilizers may be tankmixed with the Furadan spray.

RESTRICTIONS:

Do not apply within 90 days of harvest.

Do not use more than two quarts per acre per season.

Furadan spray solution must have a pH of 7 or lower. The use of a buffering agent may be necessary.

Check physical compatibility of any tank mix before making up large quantities.

Do not tank-mix with Humic Acid.

Do not use at planting.

Do not use this product through any type of irrigation system.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED.

PERSONAL PROTECTIVE EQUIPMENT AND REENTRY INTERVALS ON THIS 24C OR ON THE BASIC LABEL MUST BE FOLLOWED.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

24(c) Registrant:
FMC Corporation
Agricultural Chemical Group
1735 Market Street
Philadelphia, PA 19103

ID-920002

Code 279/2837 3/95

FMC®

Furadan and FMC - FMC Trademarks
©1995 FMC Corporation